

TRACKING YOUR TREATMENT

Use this tool to track the days and sites of your ACTIMMUNE® (Interferon gamma-1b) injections



INDICATION AND IMPORTANT SAFETY INFORMATION

What is ACTIMMUNE®

(Interferon gamma-1b) used for?

ACTIMMUNE® is part of a drug regimen used to treat Chronic Granulomatous Disease, or CGD. CGD is a genetic disorder, usually diagnosed in childhood, that affects some cells of the immune system and the body's ability to fight infections effectively. CGD is often treated (though not cured) with antibiotics, antifungals, and ACTIMMUNE.

ACTIMMUNE is also used to slow the worsening of severe, malignant osteopetrosis (SMO). SMO is a genetic disorder that affects normal bone formation and is usually diagnosed in the first few months after birth.

When should I not take ACTIMMUNE?

Don't use ACTIMMUNE if you are allergic to interferon-gamma, *E coli*-derived products, or any ingredients contained in the product.

ACTIMMUNE®
(Interferon gamma-1b)

Please review the additional Important Safety Information on page 4 and the accompanying [Full Prescribing Information](#) and [Information for the Patient/Caregiver](#) and discuss with your doctor.

Helping you get started

1

Inject 3 days a week

Pick the 3 days each week for your injection.

- Inject at the same time each day
- Don't inject 2 days in a row
- Skip at least 1 day between injections

Example schedules


Mon	Wed	Fri
1 4	3 2	5 3a
6:00 pm	5:30 pm	6:30 pm

Tue	Thu	Sat
2 5	4 1	6 3b
6:00 pm	6:30 pm	5:30 pm

DATE INJECTION SITE
 TIME OF INJECTION

2

Rotate sites for each injection



*If you select the abdomen, make sure you inject at least 2 inches away from the belly button.

3

Track your injection

On the day you inject, write the number of the injection site in the circle as shown above.

Each time you take your medication, choose an injection site that's different from the one you chose the last time. That will help make sure the skin is healed between injections.

Clinical Nurse Educator

Name: _____

Phone: _____

Specialty Pharmacy

Name: _____


Phone: _____

Please review the Important Safety Information on page 4 and the accompanying [Full Prescribing Information](#) and [Information for the Patient/Caregiver](#) and discuss with your doctor.



NAME _____

Choose a site



Track your injection

MONTH: _____

DAY	DAY	DAY
<input type="checkbox"/> ○	<input type="checkbox"/> ○	<input type="checkbox"/> ○
DAY	DAY	DAY
<input type="checkbox"/> ○	<input type="checkbox"/> ○	<input type="checkbox"/> ○
DAY	DAY	DAY
<input type="checkbox"/> ○	<input type="checkbox"/> ○	<input type="checkbox"/> ○
DAY	DAY	DAY
<input type="checkbox"/> ○	<input type="checkbox"/> ○	<input type="checkbox"/> ○

NOTES _____

Please review the Important Safety Information on page 4 and the accompanying [Full Prescribing Information](#) and [Information for the Patient/Caregiver](#) and discuss with your doctor.



IMPORTANT SAFETY INFORMATION (CONTINUED)

What warnings should I know about ACTIMMUNE?

At high doses, ACTIMMUNE can cause (flu-like) symptoms, which may worsen some pre-existing heart conditions.

ACTIMMUNE may cause decreased mental status, walking disturbances, and dizziness, particularly at very high doses. These symptoms are usually reversible within a few days upon dose reduction or discontinuation of therapy.

Bone marrow function may be suppressed with ACTIMMUNE, and decreased production of cells important to the body may occur. This effect, which can be severe, is usually reversible when the drug is discontinued or the dose is reduced.

Taking ACTIMMUNE may cause reversible changes to your liver function, particularly in patients less than 1 year old. Your doctor should monitor your liver function every 3 months, and monthly in children under 1 year.

In rare cases, ACTIMMUNE can cause severe allergic reactions and/or rash. If you experience a serious reaction to ACTIMMUNE, discontinue it immediately and contact your doctor or seek medical help.

What should I tell my healthcare provider?

Be sure to tell your doctor about all the medications you are taking.

Tell your doctor if you:

- are pregnant or plan to become pregnant or plan to nurse
- have a cardiac condition such as irregular heartbeat, heart failure, or decreased blood flow to your heart
- have a history of seizures or other neurologic disorders
- have, or have had, reduced bone marrow function. Your doctor will monitor these cells with blood tests at the beginning of therapy and at 3-month intervals on ACTIMMUNE therapy

What are the side effects of ACTIMMUNE?

The most common side effects with ACTIMMUNE are “flu-like” symptoms such as fever, headache, chills, muscle pain, or fatigue, which may decrease in severity as treatment continues. Bedtime administration of ACTIMMUNE may help reduce some of these symptoms. Acetaminophen may be helpful in preventing fever and headache.

What other medications might interact with ACTIMMUNE?

Some drugs may interact with ACTIMMUNE to potentially increase the risk of damage to your heart or nervous system, such as certain chemotherapy drugs. Tell your doctor about all other medications you are taking.

Avoid taking ACTIMMUNE at the same time as a vaccination.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

The risk information provided here is not comprehensive. To learn more, talk about ACTIMMUNE with your healthcare provider or pharmacist. The FDA-approved product labeling can be found at <http://www.ACTIMMUNE.com> or 1-866-479-6742.

Please review the accompanying Full Prescribing Information and Information for the Patient/Caregiver and discuss with your doctor.



NEED SUPPORT?

Contact your Clinical Nurse Educator for support or more information.



1 (877) 305-7704