


Way to go!

Your doctor has prescribed KRYSTEXXA® (pegloticase) and you're ready to take a big step in treating your chronic, out-of-control gout.



Here's what happens next:

KRYSTEXXA is a specialty medicine that you receive through an intravenous (IV) infusion. Specialty medicines require a few extra steps to get insurance approval and start treatment. It can take around 30 days from the time your doctor prescribes KRYSTEXXA until you get approval from your insurance provider. It can take an additional 30 days from insurance approval until you get your first infusion. Once you receive insurance approval, work with your Patient Access Liaison (PAL) to help schedule your first infusion.



KRYSTEXXA is a prescription medicine used in adults to help reduce the signs and symptoms of gout that are not controlled by other treatments.

KRYSTEXXA is not recommended if you have high levels of uric acid without a history of gout.

What is the most important information I should know about KRYSTEXXA?

Serious allergic reactions may happen in some patients who receive KRYSTEXXA.

These allergic reactions can be life threatening and usually happen within 2 hours of the infusion.

Keep reading for more details >>

Please see additional Important Safety Information for KRYSTEXXA on pages 5-7 and Medication Guide at [KRYSTEXXA.com](https://www.krystexxa.com).

KRYSTEXXA
pegloticase

PHASE 1

Prescription & enrollment



Most doctors will submit a Patient Enrollment Form to initiate enrollment into Horizon By Your Side, a patient support program. Then, they will begin the insurance approval process, inclusive of a Prior Authorization Form, which will provide important information to your insurance provider, who will begin reviewing your coverage for KRYSTEXXA.



By signing the Patient Enrollment Form, you will receive personalized support from Horizon By Your Side. You can enroll in this personalized support program by calling **1-833-469-4688, Monday–Friday 8 am–11 pm, ET**. This is an optional support program for patients, and you do not have to enroll in Horizon By Your Side to get KRYSTEXXA.



Once your enrollment is processed, you will receive a welcome call from your Patient Access Liaison (PAL). Your PAL provides you with dedicated, 1-on-1 support. They work directly with you to answer non-medical, logistical questions and provide support once you have been prescribed KRYSTEXXA and you have completed enrollment.

PHASE 2

Insurance review




The next step is the Benefits Investigation. Your insurance provider, your doctor's office or infusion center, and your Patient Access Liaison (PAL) will work together to confirm the details of your insurance coverage for KRYSTEXXA.



If you enrolled in 1-on-1 support with a PAL, they will call you to review the results of the Benefits Investigation, which is called a Summary of Benefits. It will include:

- Your insurance provider's requirements to cover KRYSTEXXA
- Financial assistance options, including what your potential out-of-pocket cost may be
- Whether your doctor needs approval from your insurance provider to prescribe KRYSTEXXA. This is called a Prior Authorization or Predetermination. Insurance providers often require a Prior Authorization for KRYSTEXXA



KRYSTEXXA is an intravenous (IV) medicine you receive through a process called an "infusion." Your doctor and/or insurance provider will work with you to find an infusion center where you will get your KRYSTEXXA treatment.




Your doctor and infusion center will work together to prepare all the necessary paperwork and submit it to your insurance provider for you. This includes the Prior Authorization, if it's needed.

PHASE 3

Insurance approval



In some cases, your insurance provider may need more information for approval. Common reasons that Prior Authorization (PA) is denied include incomplete or missing clinical information.



If you've enrolled in Horizon By Your Side, reach out to your Patient Access Liaison (PAL) if you ever have any questions about this process.



Once your insurance provider has all the information they need, they will make a coverage decision on KRYSTEXXA. If approved, you can start planning your first infusion. Your PAL is available to support and educate you through this process.

Remember: The insurance approval process may take up to 60 days, but a decision is reached sooner for many patients.



Sometimes, an insurance provider may decide not to cover KRYSTEXXA. If this happens to you, your PAL can provide options and resources for access to KRYSTEXXA.

No matter how long your insurance approval process takes, **Horizon By Your Side** is here for you, if you choose to enroll.

If you would like to enroll in Horizon By Your Side, call **1-833-469-4688**, **Monday–Friday, 8 AM–11 PM, ET.**



Important Safety Information

USE

KRYSTEXXA® (pexloticase) is a prescription medicine used in adults to help reduce the signs and symptoms of gout that are not controlled by other treatments.

KRYSTEXXA is not for use in people with too much uric acid in their bodies who do not have symptoms (asymptomatic hyperuricemia).

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about KRYSTEXXA?

Serious allergic reactions may happen in some patients who receive KRYSTEXXA. These allergic reactions can be life-threatening and usually happen within 2 hours of the infusion.

KRYSTEXXA should be given to you by a doctor or nurse in a healthcare setting where serious allergic reactions can be treated. Your doctor or nurse should watch you for any signs of a serious allergic reaction during and after your treatment with KRYSTEXXA.

Tell your doctor or nurse right away if you have any of these symptoms during or after your treatment with KRYSTEXXA:

- wheezing, shortness of breath, cough, chest tightness, chest pain, or trouble breathing
- dizziness, fainting, fast or weak heartbeat or feeling nervous
- reddening of the face, itching, hives, or feeling warm
- swelling of the throat or tongue, throat tightness, hoarse voice, or trouble swallowing

Who should not receive KRYSTEXXA?

Do not receive KRYSTEXXA if you:

- have a rare blood problem called glucose-6-phosphate dehydrogenase (G6PD) deficiency or favism. Your doctor may test you for G6PD before you start KRYSTEXXA.
- have had a serious allergic reaction to KRYSTEXXA or any of its ingredients. See the end of the Medication Guide for a complete list of ingredients in KRYSTEXXA.

Please see additional Important Safety Information for KRYSTEXXA on pages 6-7 and Medication Guide at KRYSTEXXA.com.

Important Safety Information (cont.)

What should I tell my doctor before receiving treatment with KRYSTEXXA?

Before you receive KRYSTEXXA, tell your doctor about all of your medical conditions, including if you:

- ever had any heart problems or high blood pressure.
- are pregnant or plan to become pregnant. It is not known if KRYSTEXXA will harm your unborn baby. Talk to your doctor if you are pregnant or plan to become pregnant.
- are breastfeeding or plan to breastfeed. It is not known if KRYSTEXXA passes into your breast milk. You and your doctor should decide if you will receive KRYSTEXXA or breastfeed.

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. **Do not** take any other uric acid lowering drug, such as allopurinol, febuxostat (Uloric), or probenecid, while receiving KRYSTEXXA.

KRYSTEXXA is recommended to be given with another prescription medicine called methotrexate. KRYSTEXXA may also be used alone. You and your doctor will decide the treatment that is right for you.

Prior to your treatment with KRYSTEXXA, your doctor may give you medicine to help reduce your risk of getting gout flares or an allergic reaction. Take these medicines as directed by your doctor or nurse. Your doctor will also test your uric acid levels prior to each treatment to monitor your response to KRYSTEXXA.

Your gout flares may increase in the first 3 months when you start receiving KRYSTEXXA. It's important to understand that this is happening because KRYSTEXXA is breaking down uric acid in your body. **Do not** stop receiving KRYSTEXXA even if you have a flare, as the amount of flares will decrease after 3 months of treatment. Your doctor may give you other medicines to help reduce your gout flares for the first few months after starting KRYSTEXXA.

Please see additional Important Safety Information for KRYSTEXXA on page 7 and Medication Guide at [KRYSTEXXA.com](https://www.krystexxa.com).

KRYSTEXXA
peglicase

Important Safety Information (cont.)

What are the possible side effects of KRYSTEXXA?

In KRYSTEXXA clinical trials:

The most common side effects of KRYSTEXXA when given together with methotrexate were gout flares, joint pain, coronavirus disease 2019 (COVID-19), nausea, and fatigue.

The most common side effects of KRYSTEXXA were gout flares, allergic reactions (including infusion reactions). See “What is the most important information I should know about KRYSTEXXA?”, nausea, bruising, sore throat, constipation, chest pain, coronavirus disease 2019 (COVID-19), and vomiting.

This is not a complete list of all possible side effects. Call your doctor for medical advice about side effects.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/safety/medwatch or call 1-800-FDA-1088.

For additional Important Safety Information, please see the [Medication Guide](#) and discuss with your doctor.